

Ankle Arthroscopy & Talus OCD Microfracture

INTRODUCTION

The talus is lined with articular cartilage and connects to the tibia and fibula to make up the ankle joint. An osteochondral injury occurs when the ankle joint cartilage is bruised, fractured or displaced from an injury. When cartilage injuries don't heal they often need stimulation of the underlying bone marrow to provoke a healing response. This is called a microfracture and is usually carried out with arthroscopy (keyhole surgery). Loose ankle ligaments from the time of injury often need repairing at the same time to stabilise the joint.

THE SURGERY

Ankle arthroscopy, talus OCD micro fracture and lateral ligament surgery involves a number of steps:

- General anaesthetic and IV antibiotics
- Tourniquet around the thigh
- Two small incisions (~6mm) over front of ankle
- Insertion of arthroscope (camera) and keyhole surgery instruments to clean up the cartilage injury and perform microfracture
- Incision over the outside of the ankle (~5cm)
- Reconstruction of the lateral ligaments using 2 small suture anchors inserted into the fibula bone
- Closure of incision with sutures
- Local anaesthetic block
- Surgical camboot (VACOCast)

GUIDELINES FOR POST-OP RECOVERY

HOSPITAL ADMISSION

- Most patients go home on the day of surgery

FIRST 2 WEEKS

- Elevate foot and rest
- Strictly NON-WEIGHTBEARING in surgical boot
- Mobilise with knee scooter/crutches/frame
- Boot and dressings to stay dry and intact
- Strong painkillers as required
- Aspirin 100mg and Vitamin D 5000 units per day
- Antibiotics if prescribed
- Wiggle toes to encourage circulation

2 WEEK POST-OP APPOINTMENT

- Review by nurse & removal of dressings and sutures

WEEKS 3-6

- TOUCH WEIGHTBEARING in boot
- Surgical boot on at ALL times, including in bed and showers, except for physio/exercises
- Elevate foot when not walking
- Aspirin 100mg and Vitamin D 5000 units per day
- Daily scar massage from 3 weeks post-op
- Physiotherapy review (arrange first appointment for 2-5 days after the 2-week post-op check)
- **PHYSIOTHERAPY:**
 - Commence active and passive range of movement exercises - ankle and toe flexion/extension
 - NO inversion
 - General upper body, core, hip and knee strength & conditioning

6 WEEK POST-OP APPOINTMENT

- Review by Dr Zilko

FROM 6 WEEKS

- **PHYSIOTHERAPY:**
 - Commence FULL weightbearing and transition out of boot into normal supportive shoes
 - Continue ROM and commence inversion
 - Peroneus brevis activation and strengthening
 - Gait re-training
 - Ankle stability and proprioception exercises
 - Stationary bicycle/elliptical trainer out of boot

3 MONTH POST-OP APPOINTMENT

- Review by Dr Zilko

FROM 3 MONTHS

- Return to normal activities as tolerated
- **PHYSIOTHERAPY:**
 - Straight-line jogging from 4 months and progress
 - Work-specific or sport-specific training

Full recovery is usually 6-9 months, and can take up to 18 months.

Every patient's recovery is individual and depends on the severity of the injury/disease and complexity of the surgery.