

# Ankle Fracture Fixation and Arthroscopy

## INTRODUCTION

Due to the type of your ankle fracture, you require open reduction and internal fixation. This involves restoring the normal anatomy of the bones of the ankle and holding them in place with plates and screws. Arthroscopy (keyhole surgery) is also performed to evaluate the cartilage inside the joint. Fortunately, most patients have a good result following ankle fracture surgery and are able to return to most pre-injury activities.

## THE SURGERY

Ankle fracture surgery involves a number of steps:

- General anaesthetic and IV antibiotics
- Tourniquet around the thigh
- Incision/s over fractures
- Restoration of normal anatomy (open reduction), and internal fixation with plates and screws
- Two small incisions (~6mm) over front of ankle
- Insertion of arthroscope (camera) and keyhole surgery instruments to evaluate ankle joint
- Closure of incisions with sutures
- Local anaesthetic block
- Surgical camboot (VACOCast) or plaster cast

## GUIDELINES FOR POST-OP RECOVERY

### HOSPITAL ADMISSION

- In hospital post-op for 1-2 nights
- IV antibiotics and Clexane injections

### FIRST 2 WEEKS

- Elevate foot and rest
- Strictly **NON-WEIGHTBEARING** in boot/cast
- Mobilise with knee scooter/crutches/frame
- Boot/cast and dressings to stay dry and intact
- Strong painkillers as required
- Aspirin 100mg per day
- Antibiotics for 5 days
- Wiggle toes to encourage circulation

### 2 WEEK POST-OP APPOINTMENT

- Review by nurse & removal of dressings and sutures

### WEEKS 3-6

- Surgical boot on at ALL times, including in bed, except for physio/exercises & showers (seated with shower chair)
- Elevate foot when resting
- Daily scar massage from 3 weeks post-op
- Aspirin 100mg per day
- Physiotherapy review (arrange first appointment for 2-5 days after the 2-week post-op check)
- **PHYSIOTHERAPY:**
  - Commence gentle active and passive ankle, midfoot and forefoot ROM exercises
  - Commence partial weightbearing, and progress up to full weightbearing at 6 weeks post-op if tolerated - in surgical boot for all weightbearing
  - Exercise bicycle with surgical boot on
  - General upper body, core, hip and knee exercises

### 6 WEEK POST-OP APPOINTMENT

- X-ray and review by Dr Zilko

### WEEKS 7-12

- **PHYSIOTHERAPY:**
  - Transition out of boot into normal supportive shoes
  - Gait re-training
  - Exercise bicycle +/- elliptical trainer in normal shoes
  - Continue lower limb strength and conditioning

### 12 WEEK POST-OP APPOINTMENT

- X-ray and review by Dr Zilko

### FROM 12 WEEKS

- **PHYSIOTHERAPY:**
  - Work-specific or sport-specific training
  - Safe to commence open-chain impact/running if fractures healed

Full recovery is usually 6-12 months. Some patients require removal of metalwork from 6 months following their original surgery.

*Every patient's recovery is individual and depends on the severity of the injury and complexity of the surgery.*