

Ankle Joint Fusion - Open

INTRODUCTION

Like all joints the ankle can be affected by various types of arthritis. This leads to increasing pain, swelling and stiffness, resulting in an ankle that does not function well. Ankle fusion (arthrodesis) is a very effective way to relieve the pain of ankle arthritis. Ankle fusion involves joining the two main bones of the ankle (the tibia and the talus together). After an ankle fusion most patients are able to walk without a limp and have minimal pain.

THE SURGERY

Ankle fusion surgery involves a number of steps:

- General anaesthetic, nerve block and IV antibiotics
- Tourniquet around the thigh
- Long incision over the front of the ankle joint
- Removal of any remaining cartilage
- Bone graft harvest (calcaneum or proximal tibia)
- Insertion of bone graft into the ankle joint to stimulate fusion
- Internal fixation with screws +/- a plate
- Check x-rays
- Closure of incision with sutures
- Application of plaster cast

GUIDELINES FOR POST-OP RECOVERY

HOSPITAL ADMISSION

- In hospital for 1-2 nights
- IV antibiotics and Clexane injections

FIRST 2 WEEKS

- Elevate foot and rest
- Strictly **NON-WEIGHTBEARING** in plaster cast
- Mobilise with knee scooter/crutches/frame
- Cast and dressings to stay dry and intact
- Strong painkillers as required
- Aspirin 100mg and Vitamin D 5000 units per day
- Antibiotics for 5 days
- Wiggle toes to encourage circulation

2 WEEK POST-OP APPOINTMENT

- Nurse review: removal of cast, dressings and sutures
- Change to VACOCast surgical boot

WEEKS 3-6

- Strictly **NON-WEIGHTBEARING** in boot
- Surgical boot on at **ALL** times, including in bed and showers, except to change boot liner
- Keep ankle and boot dry at all times
- Mobilise with knee scooter/crutches/frame
- Elevate foot when resting
- Aspirin 100mg and Vitamin D 5000 units per day
- Wiggle toes to encourage circulation
- Physiotherapy review (arrange first appointment for 2-5 days after the 2-week post-op check)
- **PHYSIOTHERAPY:**
 - General upper body, core, hip and knee exercises

6 WEEK POST-OP APPOINTMENT

- X-ray and review by Dr Zilko

WEEKS 7-12

- Surgical boot on at all times during the day except for physio/exercises
- Can sleep out of boot at night
- Can shower out of boot whilst on shower chair
- Elevate foot when resting
- Aspirin 100mg and Vitamin D 5000 units per day
- **PHYSIOTHERAPY:**
 - Commence partial weightbearing in surgical boot with crutches/frame, and progress up to full weightbearing as tolerated
 - Commence gentle passive and active ROM of subtalar, midfoot and forefoot joints
 - Soft tissue massage and scar mobilisation

12 WEEK POST-OP APPOINTMENT

- X-ray and review by Dr Zilko

FROM 12 WEEKS

- **PHYSIOTHERAPY:**
 - Transition out of boot into normal supportive shoes
 - Gait re-training
 - Continue lower limb strength and conditioning

Full recovery is usually 6-12 months.

Every patient's recovery is individual and depends on the severity of the injury/disease and complexity of the surgery.